

The Goldie Theorem for H -semiprime algebras [☆]

Serge Skryabin ^{a,*}, Freddy Van Oystaeyen ^b

^a Chebotarev Research Institute, Universitetskaya St. 17, 420008 Kazan, Russia

^b Department of Mathematics and Informatics, University of Antwerp, Universiteitsplein 1,
2610 Antwerp, Belgium

Received 5 September 2005

Available online 28 July 2006

Communicated by J.T. Stafford

Abstract

The main result states that, under certain assumptions about a Hopf algebra H , every H -semiprime right Noetherian H -module algebra has a quasi-Frobenius classical right quotient ring. Another question treated in the paper is concerned with the extension of H -module structures to quotient rings. These results have an application to the semiprimeness problem for smash product algebras $A \# H$.

© 2006 Elsevier Inc. All rights reserved.

Keywords: The Goldie Theorem; Quotient rings; Hopf algebras; Module algebras; H -semiprime algebras; Smash products

Introduction

Let H be a Hopf algebra over a commutative ring k . An H -module algebra A is called *H -semiprime* if A contains no nonzero nilpotent H -stable ideals. We say that A is *H -simple* if A has no nonzero proper H -stable ideals and A is *H -semisimple* if it is isomorphic to a direct product of finitely many H -simple H -module algebras. This paper is devoted to the proof of the following main result:

[☆] Both authors acknowledge support of the ESF program on Noncommutative Geometry. The first author thanks the Free University of Brussels VUB and the Mathematics Research Center of Warwick University for hospitality.

* Corresponding author.

E-mail addresses: serge.skryabin@ksu.ru (S. Skryabin), voyst@uia.ua.ac.be (F. Van Oystaeyen).